

## CONFIDENCE LIMITS FOR SMALL NUMBERS OF EVENTS IN ASTROPHYSICAL DATA

NEIL GEHRELS

Laboratory for High Energy Astrophysics, NASA/Goddard Space Flight Center

Received 1985 August 5; accepted 1985 September 30

## ABSTRACT

Convenient tables and approximate formulae are presented for confidence limits based on Poisson and binomial statistics. Poisson statistics apply when event rates are calculated from small numbers of observed events, and binomial statistics apply when ratios of two different event types are calculated from small numbers of observed events. The limits in the tables are given for all confidence levels commonly used in astrophysics.

*Subject heading:* numerical methods

## I. INTRODUCTION

In many areas of astrophysics it occasionally happens that only a small number of events of interest are detected during an observation. Examples range from the number of supernovae seen in a given period of time from a cluster of galaxies to the number of gamma rays detected during a source observation. If the goal is to determine quantities such as the event rate or the ratio of different event types, then the best approach is to repeat the measurement with a longer integration time or a larger collection factor in order to obtain enough events for an accurate measurement. In some cases, for one reason or another, this is not possible or practical, and one is forced to make the best use of the data in hand. Results are then typically quoted as upper limits at a specified confidence level or as a measured value with error bars containing a specified confidence interval. Conventionally, error bars plotted in figures are 84.13% confidence upper and lower limits containing a 68.27% confidence interval.

The calculation of limits for small numbers of counts is based on standard equations derived from Poisson and binomial statistics. Although the equations are straightforward, using them directly is cumbersome and involves interpolating in tables and executing several mathematical operations. Tables of limits for a few confidence levels have been published (Pearson and Hartley 1966; Beyer 1966), but not all levels commonly used in astrophysics are included. Also, I have found that the tables most often referenced (Beyer 1966) have inaccuracies in the last decimal place. Another shortcoming in this subject is that approximate formulae for quick estimation or for use in computer programs are not available. The purpose of this paper is to present tables of Poisson and binomial limits for all confidence levels commonly used in astrophysics and to derive easy-to-use approximate formulae for calculating the limits.

## II. CONFIDENCE LIMITS FOR POISSON STATISTICS

## a) Definitions and Numerical Solutions

We consider the case where  $n$  events are detected in a given observation. Then, based on Poisson statistics, the upper limit,  $\lambda_u$ , and lower limit,  $\lambda_l$ , of confidence level CL are defined by

$$\sum_{x=0}^n \frac{\lambda_u^x e^{-\lambda_u}}{x!} = 1 - \text{CL}, \quad (1)$$

$$\sum_{x=0}^{n-1} \frac{\lambda_l^x e^{-\lambda_l}}{x!} = \text{CL} \quad (n \neq 0) \quad (2)$$

(see, e.g., Pearson and Hartley 1966; Beyer 1966). The lower limit for  $n = 0$  is  $\lambda_l = 0.0$ . Single-sided confidence limits are defined in equations (1) and (2) and will be used throughout this paper. Double-sided intervals of level CL' can be obtained by substituting  $(1 + \text{CL}')/2$  for CL in the equations. For instance, if  $\lambda_l$  and  $\lambda_u$  are each 95% single-sided limits, then the double-sided interval  $\lambda_l$  to  $\lambda_u$  has a confidence of 90%.

In the Appendix it is shown that equations (1) and (2) are reasonable definitions of Poisson confidence limits. In particular, it is illustrated that if a large number of observers measure a real physical rate, then at least  $100 \times \text{CL}\%$  of them will assign upper limits based on their measurements that are greater than the real rate and at least  $100 \times \text{CL}\%$  will assign lower limits that are less than the real rate, for all rates. It is also shown that the upper limits are the smallest they can be and still satisfy this condition and that the lower limits the largest they can be. The limits in equations (1) and (2) therefore optimally satisfy the definition of confidence limits (see, e.g., Cramér 1945).

It is not possible to obtain exact algebraic expressions for  $\lambda_u$  and  $\lambda_l$  from equations (1) and (2). However, as shown in the next section, approximate expressions can be found that are good to a few percent. Also, for any values of  $n$  and CL, the limits can be numerically determined to any desired accuracy. Listed in Tables 1 and 2 are exact values for  $\lambda_u$  and  $\lambda_l$  at several confidence levels for  $n = 0-50$ , determined using an iterative numerical technique (Newton's method) to solve equations (1) and (2). The confidence levels that were chosen are the obvious 0.90, 0.95, 0.99, and 0.999, as well as levels corresponding to 1, 2, and 3  $\sigma$  limits for Gaussian statistics, namely 0.8413, 0.9772, and 0.9987. Also, to allow common double-sided intervals to be determined, single-sided levels of 0.975 (double-sided 0.95), 0.995 (d-s 0.99), and 0.9995 (d-s 0.999) are included.

As an example of the use of Tables 1 and 2, consider an observation where four events are detected in 10 s. The 99% confidence upper limit to the rate is then  $1.160 \text{ s}^{-1}$ , and the 99% confidence lower limit is  $0.082 \text{ s}^{-1}$ . The 99% confidence double-sided interval (CL = 0.995 column) is  $0.067-1.259 \text{ s}^{-1}$ . The measured value with error bars corresponding to 1  $\sigma$  Gaussian errors is  $0.40^{+0.32}_{-0.19} \text{ s}^{-1}$ .

TABLE 1  
POISSON SINGLE-SIDED UPPER LIMITS

$n$	CONFIDENCE LEVEL									
	0.8413 <sup>a</sup>	0.90	0.95	0.975	0.9772 <sup>a</sup>	0.99	0.995	0.9987 <sup>a</sup>	0.999	0.9995
0.....	1.841	2.303	2.996	3.689	3.783	4.605	5.298	6.608	6.908	7.601
1.....	3.300	3.890	4.744	5.572	5.683	6.638	7.430	8.900	9.233	9.999
2.....	4.638	5.322	6.296	7.225	7.348	8.406	9.274	10.87	11.23	12.05
3.....	5.918	6.681	7.754	8.767	8.902	10.05	10.98	12.68	13.06	13.93
4.....	7.163	7.994	9.154	10.24	10.39	11.60	12.59	14.39	14.79	15.71
5.....	8.382	9.275	10.51	11.67	11.82	13.11	14.15	16.03	16.45	17.41
6.....	9.584	10.53	11.84	13.06	13.22	14.57	15.66	17.62	18.06	19.05
7.....	10.77	11.77	13.15	14.42	14.59	16.00	17.13	19.17	19.63	20.65
8.....	11.95	12.99	14.43	15.76	15.94	17.40	18.58	20.69	21.16	22.22
9.....	13.11	14.21	15.71	17.08	17.27	18.78	20.00	22.18	22.66	23.75
10.....	14.27	15.41	16.96	18.39	18.58	20.14	21.40	23.64	24.13	25.26
11.....	15.42	16.60	18.21	19.68	19.87	21.49	22.78	25.08	25.59	26.74
12.....	16.56	17.78	19.44	20.96	21.16	22.82	24.14	26.51	27.03	28.20
13.....	17.70	18.96	20.67	22.23	22.43	24.14	25.50	27.91	28.45	29.65
14.....	18.83	20.13	21.89	23.49	23.70	25.45	26.84	29.31	29.85	31.08
15.....	19.96	21.29	23.10	24.74	24.95	26.74	28.16	30.69	31.24	32.50
16.....	21.08	22.45	24.30	25.98	26.20	28.03	29.48	32.06	32.62	33.90
17.....	22.20	23.61	25.50	27.22	27.44	29.31	30.79	33.42	33.99	35.29
18.....	23.32	24.76	26.69	28.45	28.68	30.58	32.09	34.76	35.35	36.68
19.....	24.44	25.90	27.88	29.67	29.90	31.85	33.38	36.10	36.70	38.05
20.....	25.55	27.05	29.06	30.89	31.13	33.10	34.67	37.44	38.04	39.41
21.....	26.66	28.18	30.24	32.10	32.34	34.35	35.95	38.76	39.37	40.76
22.....	27.76	29.32	31.41	33.31	33.55	35.60	37.22	40.07	40.70	42.11
23.....	28.87	30.45	32.59	34.51	34.76	36.84	38.48	41.38	42.02	43.45
24.....	29.97	31.58	33.75	35.71	35.96	38.08	39.74	42.69	43.33	44.78
25.....	31.07	32.71	34.92	36.90	37.16	39.31	41.00	43.98	44.64	46.11
26.....	32.16	33.84	36.08	38.10	38.36	40.53	42.25	45.28	45.94	47.42
27.....	33.26	34.96	37.23	39.28	39.55	41.76	43.50	46.56	47.23	48.74
28.....	34.35	36.08	38.39	40.47	40.74	42.98	44.74	47.84	48.52	50.04
29.....	35.45	37.20	39.54	41.65	41.92	44.19	45.98	49.12	49.80	51.35
30.....	36.54	38.32	40.69	42.83	43.10	45.40	47.21	50.39	51.08	52.64
31.....	37.63	39.43	41.84	44.00	44.28	46.61	48.44	51.66	52.36	53.94
32.....	38.72	40.54	42.98	45.17	45.46	47.81	49.67	52.92	53.63	55.23
33.....	39.80	41.65	44.13	46.34	46.63	49.01	50.89	54.18	54.90	56.51
34.....	40.89	42.76	45.27	47.51	47.80	50.21	52.11	55.43	56.16	57.79
35.....	41.97	43.87	46.40	48.68	48.97	51.41	53.32	56.69	57.42	59.06
36.....	43.06	44.98	47.54	49.84	50.14	52.60	54.54	57.93	58.67	60.34
37.....	44.14	46.08	48.68	51.00	51.30	53.79	55.75	59.18	59.93	61.60
38.....	45.22	47.19	49.81	52.16	52.46	54.98	56.96	60.42	61.17	62.87
39.....	46.30	48.29	50.94	53.31	53.62	56.16	58.16	61.66	62.42	64.13
40.....	47.38	49.39	52.07	54.47	54.78	57.35	59.36	62.89	63.66	65.39
41.....	48.46	50.49	53.20	55.62	55.93	58.53	60.56	64.13	64.90	66.64
42.....	49.53	51.59	54.32	56.77	57.09	59.71	61.76	65.36	66.14	67.90
43.....	50.61	52.69	55.45	57.92	58.24	60.88	62.96	66.58	67.37	69.14
44.....	51.68	53.78	56.57	59.07	59.39	62.06	64.15	67.81	68.60	70.39
45.....	52.76	54.88	57.69	60.21	60.54	63.23	65.34	69.03	69.83	71.63
46.....	53.83	55.97	58.82	61.36	61.69	64.40	66.53	70.25	71.06	72.88
47.....	54.90	57.07	59.94	62.50	62.83	65.57	67.72	71.47	72.28	74.11
48.....	55.98	58.16	61.05	63.64	63.97	66.74	68.90	72.68	73.51	75.35
49.....	57.05	59.25	62.17	64.78	65.12	67.90	70.08	73.90	74.72	76.58
50.....	58.12	60.34	63.29	65.92	66.26	69.07	71.27	75.11	75.94	77.81
60.....	68.79	71.20	74.39	77.23	77.60	80.62	82.99	87.11	88.01	90.01
70.....	79.41	81.99	85.40	88.44	88.83	92.06	94.58	98.96	99.91	102.0
80.....	89.98	92.73	96.35	99.57	99.98	103.4	106.1	110.7	111.7	113.9
90.....	100.5	103.4	107.2	110.6	111.1	114.7	117.4	122.3	123.3	125.7
100.....	111.0	114.1	118.1	121.6	122.1	125.8	128.8	133.8	134.9	137.4

<sup>a</sup> Corresponding to Gaussian statistics  $1\sigma = 0.8413$ ,  $2\sigma = 0.9772$ , and  $3\sigma = 0.9987$ .

TABLE 2  
POISSON SINGLE-SIDED LOWER LIMITS

<i>n</i>	CONFIDENCE LEVEL									
	0.8413	0.90	0.95	0.975	0.9772	0.99	0.995	0.9987	0.999	0.9995
1.....	0.173	0.105	5.13-2 <sup>a</sup>	2.53-2	2.30-2	1.01-2	5.01-3	1.35-3	1.00-3	5.00-4
2.....	0.708	0.532	0.355	0.242	0.230	0.149	0.103	5.29-2	4.54-2	3.20-2
3.....	1.367	1.102	0.818	0.619	0.569	0.436	0.338	0.212	0.191	0.150
4.....	2.086	1.745	1.366	1.090	1.058	0.823	0.672	0.465	0.429	0.355
5.....	2.840	2.433	1.970	1.623	1.583	1.279	1.078	0.792	0.739	0.632
6.....	3.620	3.152	2.613	2.202	2.153	1.785	1.537	1.175	1.107	0.967
7.....	4.419	3.895	3.285	2.814	2.758	2.330	2.037	1.603	1.520	1.348
8.....	5.232	4.656	3.981	3.454	3.391	2.906	2.571	2.068	1.971	1.768
9.....	6.057	5.432	4.695	4.115	4.046	3.507	3.132	2.563	2.452	2.220
10.....	6.891	6.221	5.425	4.795	4.719	4.130	3.717	3.084	2.961	2.699
11.....	7.734	7.021	6.169	5.491	5.409	4.771	4.321	3.628	3.491	3.202
12.....	8.585	7.829	6.924	6.201	6.113	5.428	4.943	4.191	4.042	3.726
13.....	9.441	8.646	7.690	6.922	6.828	6.099	5.580	4.772	4.611	4.269
14.....	10.30	9.470	8.464	7.654	7.555	6.782	6.231	5.367	5.195	4.828
15.....	11.17	10.30	9.246	8.395	8.291	7.477	6.893	5.977	5.794	5.402
16.....	12.04	11.14	10.04	9.145	9.036	8.181	7.567	6.599	6.405	5.990
17.....	12.92	11.98	10.83	9.903	9.789	8.895	8.251	7.233	7.028	6.590
18.....	13.80	12.82	11.63	10.67	10.55	9.616	8.943	7.877	7.662	7.201
19.....	14.68	13.67	12.44	11.44	11.32	10.35	9.644	8.530	8.306	7.822
20.....	15.57	14.53	13.25	12.22	12.09	11.08	10.35	9.193	8.958	8.453
21.....	16.45	15.38	14.07	13.00	12.87	11.83	11.07	9.863	9.619	9.093
22.....	17.35	16.24	14.89	13.79	13.65	12.57	11.79	10.54	10.29	9.741
23.....	18.24	17.11	15.72	14.58	14.44	13.33	12.52	11.23	10.96	10.40
24.....	19.14	17.97	16.55	15.38	15.23	14.09	13.26	11.92	11.65	11.06
25.....	20.03	18.84	17.38	16.18	16.03	14.85	14.00	12.62	12.34	11.73
26.....	20.93	19.72	18.22	16.98	16.83	15.62	14.74	13.32	13.03	12.41
27.....	21.84	20.59	19.06	17.79	17.64	16.40	15.49	14.03	13.73	13.09
28.....	22.74	21.47	19.90	18.61	18.45	17.17	16.25	14.75	14.44	13.78
29.....	23.65	22.35	20.75	19.42	19.26	17.96	17.00	15.47	15.15	14.47
30.....	24.55	23.23	21.59	20.24	20.07	18.74	17.77	16.19	15.87	15.17
31.....	25.46	24.11	22.44	21.06	20.89	19.53	18.53	16.92	16.59	15.87
32.....	26.37	25.00	23.30	21.89	21.71	20.32	19.30	17.65	17.32	16.58
33.....	27.28	25.89	24.15	22.72	22.54	21.12	20.08	18.39	18.05	17.30
34.....	28.20	26.77	25.01	23.55	23.36	21.92	20.86	19.13	18.78	18.01
35.....	29.11	27.66	25.87	24.38	24.19	22.72	21.64	19.88	19.52	18.73
36.....	30.03	28.56	26.73	25.21	25.03	23.53	22.42	20.63	20.26	19.46
37.....	30.94	29.45	27.59	26.05	25.86	24.33	23.21	21.38	21.00	20.19
38.....	31.86	30.34	28.46	26.89	26.70	25.14	24.00	22.14	21.75	20.92
39.....	32.78	31.24	29.33	27.73	27.53	25.96	24.79	22.89	22.51	21.66
40.....	33.70	32.14	30.20	28.58	28.38	26.77	25.59	23.66	23.26	22.40
41.....	34.62	33.04	31.07	29.42	29.22	27.59	26.38	24.42	24.02	23.14
42.....	35.55	33.94	31.94	30.27	30.06	28.41	27.18	25.19	24.78	23.88
43.....	36.47	34.84	32.81	31.12	30.91	29.23	27.99	25.96	25.54	24.63
44.....	37.39	35.74	33.69	31.97	31.76	30.05	28.79	26.73	26.31	25.38
45.....	38.32	36.65	34.56	32.82	32.61	30.88	29.60	27.51	27.08	26.14
46.....	39.24	37.55	35.44	33.68	33.46	31.70	30.41	28.29	27.85	26.89
47.....	40.17	38.46	36.32	34.53	34.31	32.53	31.22	29.07	28.62	27.65
48.....	41.10	39.36	37.20	35.39	35.17	33.36	32.03	29.85	29.40	28.42
49.....	42.02	40.27	38.08	36.25	36.02	34.20	32.85	30.63	30.18	29.18
50.....	42.95	41.18	38.96	37.11	36.88	35.03	33.66	31.42	30.96	29.95
60.....	52.28	50.31	47.85	45.79	45.53	43.46	41.93	39.40	38.88	37.73
70.....	61.65	59.51	56.83	54.57	54.29	52.02	50.33	47.54	46.96	45.70
80.....	71.07	68.77	65.88	63.44	63.13	60.67	58.84	55.81	55.18	53.80
90.....	80.53	78.08	74.98	72.37	72.04	69.41	67.44	64.18	63.51	62.02
100.....	90.02	87.42	84.14	81.36	81.02	78.22	76.12	72.65	71.92	70.33

<sup>a</sup> 5.13-2 means  $5.13 \times 10^{-2}$ .

b) Approximate Algebraic Expressions

The standard method for determining Poisson confidence limits makes use of the relation between the Poisson sum and the  $\chi^2$  probability function

$$\sum_{x=0}^{n-1} \frac{\lambda^x e^{-\lambda}}{x!} = 1 - P(\chi^2|v) \tag{3}$$

(see, e.g., Abramowitz and Stegun 1965), where  $P(\chi^2|v)$  is the  $\chi^2$  probability for  $v$  degrees of freedom with  $\chi^2 = 2\lambda$  and  $v = 2n$ . From equation (3), the Poisson upper and lower limits of confidence CL defined by equations (1) and (2) are related to the  $\chi^2$  probability function by

$$P(2\lambda_u|2n + 2) = CL \tag{4}$$

$$P(2\lambda_l|2n) = 1 - CL \tag{5}$$

Thus,  $\lambda_u$  and  $\lambda_l$  can be determined using tables of percentage points of the  $\chi^2$  distribution. This method is straightforward but is definitely less convenient than tables of the limits themselves. Also, since no  $\chi^2$  table covers all confidence levels commonly used, interpolations and extrapolations are often required. Equations (4) and (5) are, however, good starting points for developing approximate algebraic expressions for the limits.

i) Upper Limits

Several approximate inverse relations giving  $\chi^2$  in terms of  $P(\chi^2|v)$  are listed by Abramowitz and Stegun (1965). In this section, these relations will be used to obtain several approximate expressions for  $\lambda_u$ . Their accuracy will then be compared and a recommended numerical procedure given. All of the equations in Abramowitz and Stegun make use of the variable  $y_a$  defined as the Gaussian integral value with the same probability as  $P(\chi^2|v)$ ; i.e., if  $P(\chi_a^2|v) = a$ , then  $y_a$  is such that  $G(y_a) = (2\pi)^{-1/2} \int_{-\infty}^{y_a} e^{-t^2/2} dt = a$ . In our final equations we use the variable  $S = |y_a|$ , which is the equivalent Gaussian number of  $\sigma$  corresponding to the confidence level. Values of  $S$  for the common confidence levels are listed in Table 3, along with other data pertaining to the next section.

The first approximate relation is

$$\chi_a^2 \approx \frac{1}{2}(y_a + \sqrt{2v - 1})^2, \tag{6}$$

which, combined with equation (4), gives

$$\lambda_u \approx n + S \sqrt{n + \frac{3}{4}} + \frac{S^2 + 3}{4} \tag{7}$$

Another approximate relation is

$$\chi_a^2 \approx v \left( 1 - \frac{2}{9v} + y_a \sqrt{\frac{2}{9v}} \right)^3, \tag{8}$$

which gives

$$\lambda_u \approx (n + 1) \left[ 1 - \frac{1}{9(n + 1)} + \frac{S}{3\sqrt{n + 1}} \right]^3 \tag{9}$$

A final approximate expression is obtained by expanding equation (9) and keeping only terms of order  $(n + 1)^\alpha$  where  $\alpha \geq 0$ , giving

$$\lambda_u \approx n + S\sqrt{n + 1} + \frac{S^2 + 2}{3} \tag{10}$$

In Table 4 we compare the accuracy of equations (7), (9), and (10) for several values of  $n$  at the two extreme confidence levels discussed in this paper. The table lists the exact values from Table 1 in the column under  $\lambda_u$  and gives the values from the approximations along with (in parentheses) the percentage error with respect to  $\lambda_u$ . The Gaussian limit  $n + S(n)^{1/2}$  is also given for comparison. Equation (7) is quite good for  $CL = 0.8413$  but becomes poor at higher confidence levels. Equation (9) is the most complicated of the approximate expressions but is best overall—within 3% of the true limit for all values of  $n$  and all confidence levels considered in this paper. Equation (10) is fair at low confidence levels but becomes very accurate at high confidence levels where it actually does better than equation (9) from which it was derived. Its accuracy is better than 10% for all values of  $n$  for all confidence levels and better than 1% for  $n \geq 4$ .

*Recommended approximation:* For applications where upper limits good to within 10% are accurate enough, use the simple expression in equation (10),  $\lambda_u \approx n + S(n + 1)^{1/2} + (S^2 + 2)/3$ . If higher accuracy is required, use the exact upper limits for the smallest few  $n$  and equation (10) for all others or use the more complicated equation (9). For the special case of  $S = 1$ , the simple expression in equation (7),  $\lambda_u \approx n + (n + \frac{3}{4})^{1/2} + 1$ , is good to better than 1.5% for all  $n$ . The Gaussian statistics limit  $\lambda_u \approx n + S(n)^{1/2}$  becomes accurate at the few percent level only for  $n > 100$ .

ii) Lower Limits

To obtain approximate expressions for the Poisson lower limits, the same approximate  $\chi^2$  inverse relations from the last section are used. In this case, the starting point is the relationship between the Poisson confidence lower limit and the  $\chi^2$  probability function expressed in equation (5). Combining equation (5) with equation (6) gives

$$\lambda_l \approx n - S \sqrt{n - \frac{1}{4}} + \frac{S^2 - 1}{4}, \tag{11}$$

and with equation (8) gives

$$\lambda_l \approx n \left( 1 - \frac{1}{9n} - \frac{S}{3\sqrt{n}} \right)^3 \tag{12}$$

TABLE 3

DATA FOR APPROXIMATE EXPRESSIONS FOR POISSON LIMITS

PARAMETER	CONFIDENCE LEVEL									
	0.8413	0.900	0.950	0.975	0.9772	0.990	0.995	0.9987	0.999	0.9995
$S^a$ .....	1.000	1.282	1.645	1.960	2.000	2.326	2.576	3.000	3.090	3.291
$\beta$ .....	0.0	0.010	0.031	0.058	0.062	0.103	0.141	0.222	0.241	0.287
$\gamma$ .....	...	-4.00	-2.50	-2.22	-2.19	-2.07	-2.00	-1.88	-1.85	-1.80

<sup>a</sup> Number of Gaussian  $\sigma$ .

TABLE 4  
COMPARISON OF POISSON UPPER LIMIT APPROXIMATE EXPRESSIONS

n	CL	S	$\lambda_u$	GAUSSIAN LIMIT		EQUATION (7)		EQUATION (9)		EQUATION (10)	
				Value	(% Error <sup>a</sup> )	Value	(% Error)	Value	(% Error)	Value	(% Error)
0.....	0.8413	1.0	1.841	0.000	(-)	1.866	(1.4)	1.826	(0.8)	2.000	(8.6)
1.....	0.8413	1.0	3.300	2.000	(39)	3.323	(0.7)	3.287	(0.4)	3.414	(3.5)
2.....	0.8413	1.0	4.638	3.414	(26)	4.658	(0.4)	4.627	(0.2)	4.732	(2.0)
3.....	0.8413	1.0	5.918	4.732	(20)	5.936	(0.3)	5.909	(0.2)	6.000	(1.4)
4.....	0.8413	1.0	7.163	6.000	(16)	7.179	(0.2)	7.154	(0.1)	7.236	(1.0)
10.....	0.8413	1.0	14.27	13.16	(7.7)	14.28	(0.08)	14.26	(0.04)	14.32	(0.3)
100.....	0.8413	1.0	111.0	110.0	(0.9)	111.0	(<) <sup>b</sup>	111.0	(<)	111.0	(0.01)
0.....	0.9995	3.291	7.601	0.000	(-)	6.308	(17)	7.832	(3.0)	7.568	(0.4)
1.....	0.9995	3.291	9.999	4.291	(57)	8.811	(12)	10.18	(1.8)	9.931	(0.7)
2.....	0.9995	3.291	12.05	6.654	(45)	10.92	(9.4)	12.20	(1.3)	11.98	(0.6)
3.....	0.9995	3.291	13.93	8.700	(38)	12.83	(7.9)	14.07	(1.0)	13.86	(0.5)
4.....	0.9995	3.291	15.71	10.58	(33)	14.63	(6.9)	15.83	(0.8)	15.64	(0.5)
10.....	0.9995	3.291	25.26	20.41	(19)	24.25	(4.0)	25.34	(0.3)	25.19	(0.3)
100.....	0.9995	3.291	137.4	132.9	(3.3)	136.5	(0.6)	137.4	(0.02)	137.4	(0.02)

<sup>a</sup> % error = 100 × |value -  $\lambda_u$ | /  $\lambda_u$ .

<sup>b</sup> Symbol "<" means less than 0.01%.

Expanding equation (12) and keeping only terms of order  $n^\alpha$ , where  $\alpha \geq 0$ , results in

$$\lambda_l \approx n - S\sqrt{n} + \frac{S^2 - 1}{3}. \tag{13}$$

As will be shown below, none of the above three equations are adequate approximations for all confidence levels discussed in this paper. I have therefore devised a slightly more complex expression based on equation (12) that is quite accurate for all

$n$  and for all confidence levels in Table 3. The equation is

$$\lambda_l \approx n \left( 1 - \frac{1}{9n} - \frac{S}{3\sqrt{n}} + \beta n^\gamma \right)^3, \tag{14}$$

where  $\beta$  and  $\gamma$  are free parameters that are adjusted for each confidence level. The best values are listed in Table 3 and shown graphically in Figure 1.

In Table 5 the accuracy of equations (11), (12), (13), and (14) are compared. Equations (11) and (12) are possibilities for




FIG. 1.—Parameters  $\beta$  and  $\gamma$  from Poisson lower limit approximation, eq. (14). Dots represent values in Table 3.

TABLE 5  
COMPARISON OF POISSON LOWER LIMIT APPROXIMATE EXPRESSIONS

n	CL	S	$\lambda_i$	GAUSSIAN LIMIT		EQUATION (11)		EQUATION (12)		EQUATION (13)		EQUATION (14)	
				Value	(% Error <sup>a</sup> )	Value	(% Error)	Value	(% Error)	Value	(% error)	Value	(% Error)
1.....	0.8413	1.0	0.173	0.000	(-)	0.134	(22)	0.171	(0.7)	0.000	(-)	0.171	(0.7)
2.....	0.8413	1.0	0.708	0.586	(17)	0.677	(4.4)	0.712	(0.5)	0.586	(17)	0.712	(0.5)
3.....	0.8413	1.0	1.367	1.268	(7.3)	1.342	(1.9)	1.372	(0.4)	1.268	(7.3)	1.372	(0.4)
4.....	0.8413	1.0	2.086	2.000	(4.1)	2.064	(1.1)	2.091	(0.3)	2.000	(4.1)	2.091	(0.3)
10.....	0.8413	1.0	6.891	6.838	(0.8)	6.878	(0.2)	6.896	(0.07)	6.838	(0.8)	6.896	(0.07)
100.....	0.8413	1.0	90.02	90.00	(0.02)	90.01	(< <sup>b</sup> )	90.02	(<)	90.00	(0.02)	90.02	(<)
1.....	0.9995	3.291	5.00-4	-2.29	(-)	0.608	(-)	-0.009	(-)	0.986	(-)	4.91-4	(1.8)
2.....	0.9995	3.291	3.20-2	-2.65	(-)	0.104	(-)	0.010	(70)	0.623	(-)	3.17-2	(0.9)
3.....	0.9995	3.291	0.150	-2.70	(-)	1.64-4	(-)	0.107	(28)	0.577	(-)	0.151	(1.0)
4.....	0.9995	3.291	0.355	-2.58	(-)	0.085	(76)	0.304	(14)	0.695	(-)	0.358	(0.8)
10.....	0.9995	3.291	2.699	-0.41	(-)	2.182	(19)	2.646	(2.0)	2.870	(6.3)	2.703	(0.1)
100.....	0.9995	3.291	70.33	67.09	(4.6)	69.59	(1.1)	70.30	(0.01)	70.37	(0.05)	70.32	(0.01)

<sup>a</sup> % error =  $100 \times |\text{value} - \lambda_i|/\lambda_i$ .

<sup>b</sup> Symbol "<" means less than 0.01%.

CL = 0.8413 but fail for higher confidence levels. Equation (13) is poor for all cases. Equation (14) is the most complex equation but is accurate to better than 2% for all values of  $n$  and all confidence levels considered in this paper. For  $n \geq 10$  it is accurate to better than  $\sim 0.1\%$ .

*Recommended approximation:* For all cases use equation (14),  $\lambda_i \approx n(1 - 1/9n - S/3(n)^{1/2} + \beta n^\gamma)^3$ , where  $\beta$  and  $\gamma$  are given in Table 3 ( $\beta = 0$  for the special case of  $S = 1$ ). The results are accurate to better than 2%. For the general  $S$ , the Gaussian statistics limit  $\lambda_i \approx n - S(n)^{1/2}$  becomes accurate at the few percent level only for  $n > 100$ , although for  $S = 1$  it is accurate to better than 1% for  $n \geq 8$ .

### III. CONFIDENCE LIMITS FOR BINOMIAL STATISTICS

#### a) Relation to Abundance Measurements

We consider next the case where an observer is measuring two different kinds of distinguishable events. It is assumed that the source of each event type is random, so that the number of events of each type detected in a given observation time is distributed according to Poisson statistics. The objective is to obtain confidence limits on the ratio of the two event rates based on the measurement of a small number of events. Examples of this kind of study are elemental and isotopic abundance ratios in cosmic-ray astrophysics, spectral-line ratios in X-ray and  $\gamma$ -ray astronomy, and supernova type ratios in stellar astronomy.

The joint probability function for the two event types can be related to a combination of Poisson and binomial distributions. If  $n_1$  and  $n_2$  are the number of observed events of types 1 and 2, distributed according to Poisson statistics with true rates of  $\lambda_1$  and  $\lambda_2$ , then the joint probability  $f(n_1, n_2)$  of observing  $n_1$  and  $n_2$  is

$$f(n_1, n_2) = \frac{\lambda_1^{n_1} e^{-\lambda_1}}{n_1!} \frac{\lambda_2^{n_2} e^{-\lambda_2}}{n_2!} = \frac{(\lambda_1 + \lambda_2)^{n_1+n_2} e^{-(\lambda_1 + \lambda_2)}}{(n_1 + n_2)!} \binom{n_1 + n_2}{n_1} p_1^{n_1} (1 - p_1)^{n_2}, \quad (15)$$

where  $p_1 = \lambda_1/(\lambda_1 + \lambda_2)$  and  $\binom{n_1+n_2}{n_1}$  is the binomial coefficient  $(n_1 + n_2)!/n_1! n_2!$ . Thus, the joint probability is equal to the

Poisson probability for the combined rate times the binomial probability for obtaining specifically  $n_1$  and  $n_2$  given that the combined number of events observed was  $n_1 + n_2$ . Another way of viewing this is as follows: a source randomly emitting two types of events, each with its own average rate, is equivalent to a source randomly emitting events of unknown type at an average rate equal to the sum of the individual rates, with a weighted coin flipped for each event to determine its type.

The goal in the following sections will be to find confidence limits for  $p_1$  in equation (15) or equivalently for the ratio of event rates,  $r = \lambda_1/\lambda_2 = p_1/(1 - p_1)$ . Binomial statistics will be used since, as shown above, they determine the event ratios.

#### b) Definitions and Numerical Solutions

For a binomial distribution, the upper limit,  $p_{1u}$ , and lower limit,  $p_{1l}$ , for the ratio of type 1 events to total events (type 1 plus type 2) are defined by

$$\sum_{x=0}^{n_1} \binom{n}{x} p_{1u}^x (1 - p_{1u})^{n-x} = 1 - \text{CL} \quad (n_1 \neq n), \quad (16)$$

$$\sum_{x=0}^{n_1-1} \binom{n}{x} p_{1l}^x (1 - p_{1l})^{n-x} = \text{CL} \quad (n_1 \neq 0) \quad (17)$$

(see, e.g., Pearson and Hartley 1966; Beyer 1966), where CL is the single-sided confidence level of the limits and  $n = n_1 + n_2$ . For  $n_1 = n$  ( $n_2 = 0$ )  $p_{1u} = 1.0$  and for  $n_1 = 0$   $p_{1l} = 0.0$ .

Equations (16) and (17) were derived for measurements with fixed  $n$ , which is not the case for ratios computed from two random rates. For two random rates that are both small, these equations give somewhat conservative limits. As an example, for all cases with both true rates less than 5.0, the fraction of observers with upper limits greater than the true ratio is always more than 0.98 for CL = 0.95. The fraction does approach 0.95 for large rates, and the limits defined here do satisfy the definition of confidence limits (see Appendix) for all rates and confidence levels. As far as I know, these limits are the best one can do for abundance measurements.

As was the case for the Poisson limits, exact general algebraic expressions for  $p_{1u}$  and  $p_{1l}$  can not be obtained from equations (16) and (17). However, as will be discussed below, there are exact solutions for some special cases, and general approximate algebraic expressions can be derived. Also, for

TABLE 6  
BINOMIAL SINGLE-SIDED UPPER LIMITS<sup>a</sup>

$n_1$	$n_2$	CONFIDENCE LEVEL									
		0.8413	0.90	0.95	0.975	0.9772	0.99	0.995	0.9987	0.999	0.9995
0.....	1	0.841	0.9000	0.9500	0.9750	0.9772	0.9 <sup>2</sup> 00 <sup>b</sup>	0.9 <sup>2</sup> 50	0.9 <sup>2</sup> 87	0.9 <sup>3</sup> 00	0.9 <sup>3</sup> 50
	2	0.602	0.684	0.776	0.842	0.849	0.9000	0.9293	0.9633	0.9684	0.9776
	3	0.459	0.536	0.632	0.708	0.717	0.785	0.829	0.889	0.9000	0.9206
	4	0.369	0.438	0.527	0.602	0.612	0.684	0.734	0.808	0.822	0.850
	5	0.308	0.369	0.451	0.522	0.531	0.602	0.653	0.733	0.749	0.781
	6	0.264	0.319	0.393	0.459	0.468	0.536	0.586	0.668	0.684	0.718
	7	0.231	0.280	0.348	0.410	0.418	0.482	0.531	0.611	0.627	0.662
	8	0.206	0.250	0.312	0.369	0.377	0.438	0.484	0.562	0.578	0.613
	9	0.185	0.226	0.283	0.336	0.343	0.401	0.445	0.520	0.536	0.570
	10	0.168	0.206	0.259	0.308	0.315	0.369	0.411	0.484	0.499	0.532
	100	0.0182	0.0228	0.0295	0.0362	0.0371	0.0450	0.0516	0.0639	0.0667	0.0732
1.....	1	0.9172	0.9487	0.9747	0.9874	0.9886	0.9 <sup>2</sup> 50	0.9 <sup>2</sup> 75	0.9 <sup>3</sup> 32	0.9 <sup>3</sup> 50	0.9 <sup>3</sup> 75
	2	0.748	0.804	0.865	0.9057	0.9102	0.9411	0.9586	0.9786	0.9816	0.9870
	3	0.618	0.680	0.751	0.806	0.812	0.859	0.889	0.9291	0.9360	0.9493
	4	0.524	0.584	0.657	0.716	0.724	0.778	0.815	0.868	0.878	0.898
	5	0.454	0.510	0.582	0.641	0.649	0.706	0.746	0.807	0.819	0.843
	6	0.400	0.453	0.521	0.579	0.586	0.643	0.685	0.750	0.763	0.789
	7	0.357	0.406	0.471	0.527	0.534	0.590	0.632	0.698	0.711	0.740
	8	0.323	0.368	0.429	0.482	0.489	0.544	0.585	0.652	0.665	0.694
	9	0.294	0.337	0.394	0.445	0.452	0.504	0.544	0.610	0.624	0.653
	10	0.270	0.310	0.364	0.413	0.419	0.470	0.509	0.573	0.587	0.616
	100	0.0323	0.0380	0.0461	0.0539	0.0550	0.0639	0.0713	0.0848	0.0878	0.0947
2.....	1	0.9440	0.9655	0.9830	0.9 <sup>2</sup> 16	0.9 <sup>2</sup> 24	0.9 <sup>2</sup> 67	0.9 <sup>2</sup> 83	0.9 <sup>3</sup> 55	0.9 <sup>3</sup> 67	0.9 <sup>3</sup> 83
	2	0.815	0.857	0.9024	0.9324	0.9357	0.9580	0.9706	0.9848	0.9870	0.9 <sup>2</sup> 08
	3	0.703	0.753	0.811	0.853	0.858	0.894	0.9172	0.9473	0.9524	0.9625
	4	0.615	0.667	0.729	0.777	0.783	0.827	0.856	0.898	0.9060	0.9215
	5	0.546	0.596	0.659	0.710	0.716	0.764	0.797	0.847	0.856	0.876
	6	0.490	0.538	0.600	0.651	0.657	0.707	0.742	0.797	0.807	0.830
	7	0.444	0.490	0.550	0.600	0.606	0.656	0.693	0.750	0.761	0.785
	8	0.405	0.450	0.507	0.556	0.562	0.612	0.648	0.707	0.718	0.744
	9	0.373	0.415	0.470	0.518	0.524	0.572	0.608	0.667	0.679	0.705
	10	0.346	0.386	0.438	0.484	0.490	0.537	0.573	0.632	0.644	0.670
	100	0.0449	0.0513	0.0604	0.0690	0.0702	0.0799	0.0877	0.102	0.105	0.112
3.....	1	0.9577	0.9740	0.9873	0.9 <sup>2</sup> 37	0.9 <sup>2</sup> 43	0.9 <sup>2</sup> 75	0.9 <sup>2</sup> 87	0.9 <sup>3</sup> 66	0.9 <sup>3</sup> 75	0.9 <sup>3</sup> 87
	2	0.853	0.888	0.9236	0.9473	0.9498	0.9673	0.9771	0.9882	0.9899	0.9 <sup>2</sup> 29
	3	0.757	0.799	0.847	0.882	0.886	0.9153	0.9337	0.9580	0.9621	0.9701
	4	0.676	0.721	0.775	0.816	0.821	0.858	0.882	0.9170	0.9233	0.9361
	5	0.610	0.655	0.711	0.755	0.761	0.802	0.830	0.872	0.880	0.897
	6	0.555	0.599	0.655	0.701	0.706	0.750	0.781	0.828	0.837	0.856
	7	0.508	0.552	0.607	0.652	0.658	0.703	0.735	0.785	0.795	0.816
	8	0.469	0.511	0.564	0.610	0.615	0.660	0.693	0.745	0.756	0.778
	9	0.435	0.475	0.527	0.572	0.577	0.622	0.655	0.708	0.719	0.742
	10	0.405	0.444	0.495	0.538	0.544	0.588	0.621	0.674	0.685	0.709
	100	0.0566	0.0637	0.0736	0.0828	0.0840	0.0942	0.103	0.117	0.121	0.128
4.....	1	0.9660	0.9791	0.9898	0.9 <sup>2</sup> 49	0.9 <sup>2</sup> 54	0.9 <sup>2</sup> 80	0.9 <sup>3</sup> 00	0.9 <sup>3</sup> 73	0.9 <sup>3</sup> 80	0.9 <sup>4</sup> 00
	2	0.879	0.9074	0.9372	0.9567	0.9588	0.9732	0.9813	0.9 <sup>2</sup> 04	0.9 <sup>2</sup> 17	0.9 <sup>2</sup> 42
	3	0.794	0.830	0.871	0.9010	0.9044	0.9292	0.9447	0.9650	0.9684	0.9751
	4	0.720	0.760	0.807	0.843	0.847	0.879	0.9001	0.9298	0.9352	0.9460
	5	0.658	0.699	0.749	0.788	0.793	0.829	0.854	0.891	0.897	0.9117
	6	0.605	0.646	0.696	0.738	0.743	0.782	0.809	0.851	0.859	0.875
	7	0.559	0.599	0.650	0.692	0.697	0.738	0.767	0.812	0.821	0.839
	8	0.519	0.559	0.609	0.651	0.656	0.698	0.728	0.775	0.784	0.804
	9	0.485	0.523	0.573	0.614	0.619	0.661	0.691	0.740	0.750	0.771
	10	0.455	0.492	0.540	0.581	0.586	0.627	0.658	0.707	0.717	0.739
	100	0.0678	0.0754	0.0859	0.0956	0.0968	0.108	0.116	0.132	0.135	0.143
5.....	1	0.9716	0.9826	0.9 <sup>2</sup> 15	0.9 <sup>2</sup> 58	0.9 <sup>2</sup> 62	0.9 <sup>2</sup> 83	0.9 <sup>3</sup> 16	0.9 <sup>3</sup> 77	0.9 <sup>3</sup> 83	0.9 <sup>4</sup> 17
	2	0.896	0.9212	0.9466	0.9633	0.9651	0.9773	0.9842	0.9 <sup>2</sup> 19	0.9 <sup>2</sup> 30	0.9 <sup>2</sup> 51
	3	0.821	0.853	0.889	0.9148	0.9177	0.9392	0.9525	0.9700	0.9730	0.9787
	4	0.754	0.790	0.831	0.863	0.867	0.895	0.9132	0.9391	0.9438	0.9532
	5	0.695	0.733	0.778	0.813	0.817	0.850	0.872	0.9041	0.9102	0.9227
	6	0.644	0.682	0.729	0.766	0.771	0.806	0.831	0.868	0.875	0.890
	7	0.600	0.638	0.685	0.723	0.728	0.765	0.791	0.832	0.840	0.857
	8	0.561	0.598	0.645	0.684	0.689	0.727	0.755	0.798	0.806	0.824
	9	0.526	0.563	0.610	0.649	0.653	0.692	0.720	0.765	0.774	0.793
	10	0.496	0.532	0.577	0.616	0.621	0.660	0.688	0.734	0.743	0.763
	100	0.0785	0.0865	0.0975	0.108	0.109	0.120	0.129	0.145	0.148	0.156

## CONFIDENCE LIMITS

TABLE 6—Continued

$n_1$	$n_2$	CONFIDENCE LEVEL									
		0.8413	0.90	0.95	0.975	0.9772	0.99	0.995	0.9987	0.999	0.9995
6.....	1	0.9756	0.9851	0.9 <sup>2</sup> 27	0.9 <sup>2</sup> 64	0.9 <sup>2</sup> 67	0.9 <sup>2</sup> 86	0.9 <sup>3</sup> 28	0.9 <sup>3</sup> 81	0.9 <sup>3</sup> 86	0.9 <sup>4</sup> 29
	2	0.9097	0.9314	0.9536	0.9681	0.9697	0.9803	0.9863	0.9 <sup>2</sup> 30	0.9 <sup>2</sup> 40	0.9 <sup>2</sup> 57
	3	0.842	0.871	0.9023	0.9251	0.9278	0.9467	0.9584	0.9737	0.9763	0.9814
	4	0.780	0.812	0.850	0.878	0.882	0.9068	0.9232	0.9462	0.9504	0.9587
	5	0.725	0.759	0.800	0.833	0.836	0.866	0.886	0.9146	0.9201	0.9312
	100	0.531	0.565	0.609	0.646	0.650	0.687	0.713	0.756	0.764	0.783
7.....	1	0.9786	0.9869	0.9 <sup>2</sup> 36	0.9 <sup>2</sup> 68	0.9 <sup>2</sup> 71	0.9 <sup>2</sup> 87	0.9 <sup>3</sup> 37	0.9 <sup>3</sup> 83	0.9 <sup>3</sup> 87	0.9 <sup>4</sup> 37
	2	0.9199	0.9392	0.9590	0.9719	0.9732	0.9826	0.9879	0.9 <sup>2</sup> 38	0.9 <sup>2</sup> 47	0.9 <sup>2</sup> 62
	3	0.858	0.884	0.9127	0.9333	0.9356	0.9525	0.9630	0.9767	0.9790	0.9834
	4	0.801	0.831	0.865	0.891	0.894	0.9163	0.9312	0.9518	0.9556	0.9630
	5	0.750	0.781	0.819	0.848	0.852	0.879	0.897	0.9230	0.9279	0.9380
	100	0.562	0.594	0.636	0.671	0.675	0.709	0.734	0.774	0.782	0.800
8.....	1	0.9810	0.9884	0.9 <sup>2</sup> 43	0.9 <sup>2</sup> 72	0.9 <sup>2</sup> 74	0.9 <sup>2</sup> 89	0.9 <sup>3</sup> 44	0.9 <sup>3</sup> 85	0.9 <sup>3</sup> 89	0.9 <sup>4</sup> 44
	2	0.9280	0.9455	0.9632	0.9748	0.9760	0.9845	0.9891	0.9 <sup>2</sup> 44	0.9 <sup>2</sup> 52	0.9 <sup>2</sup> 66
	3	0.872	0.895	0.9212	0.9398	0.9419	0.9572	0.9667	0.9790	0.9811	0.9851
	4	0.819	0.846	0.877	0.9008	0.9035	0.9241	0.9376	0.9564	0.9598	0.9665
	5	0.770	0.799	0.834	0.861	0.865	0.889	0.9058	0.9299	0.9344	0.9436
	100	0.588	0.620	0.659	0.692	0.697	0.729	0.753	0.790	0.798	0.814
9.....	1	0.9829	0.9895	0.9 <sup>2</sup> 49	0.9 <sup>2</sup> 75	0.9 <sup>2</sup> 77	0.9 <sup>3</sup> 00	0.9 <sup>3</sup> 50	0.9 <sup>3</sup> 86	0.9 <sup>4</sup> 00	0.9 <sup>4</sup> 50
	2	0.9347	0.9505	0.9667	0.9772	0.9783	0.9859	0.9 <sup>2</sup> 02	0.9 <sup>2</sup> 50	0.9 <sup>2</sup> 57	0.9 <sup>2</sup> 70
	3	0.883	0.9043	0.9281	0.9451	0.9471	0.9610	0.9697	0.9809	0.9828	0.9864
	4	0.833	0.858	0.887	0.9091	0.9116	0.9305	0.9429	0.9601	0.9632	0.9694
	5	0.787	0.815	0.847	0.872	0.875	0.898	0.9134	0.9356	0.9398	0.9483
	100	0.612	0.642	0.680	0.711	0.715	0.746	0.768	0.804	0.811	0.826
10.....	1	0.9844	0.9 <sup>2</sup> 05	0.9 <sup>2</sup> 53	0.9 <sup>2</sup> 77	0.9 <sup>2</sup> 79	0.9 <sup>3</sup> 09	0.9 <sup>3</sup> 54	0.9 <sup>3</sup> 88	0.9 <sup>4</sup> 09	0.9 <sup>4</sup> 55
	2	0.9402	0.9548	0.9695	0.9791	0.9802	0.9872	0.9 <sup>2</sup> 10	0.9 <sup>2</sup> 54	0.9 <sup>2</sup> 61	0.9 <sup>2</sup> 72
	3	0.892	0.9120	0.9340	0.9496	0.9514	0.9642	0.9722	0.9825	0.9842	0.9876
	4	0.846	0.869	0.896	0.9161	0.9185	0.9360	0.9474	0.9633	0.9662	0.9719
	5	0.802	0.828	0.858	0.882	0.885	0.9056	0.9199	0.9405	0.9444	0.9522
	6	0.762	0.790	0.822	0.848	0.851	0.875	0.891	0.9159	0.9206	0.9302
	7	0.726	0.754	0.788	0.816	0.819	0.845	0.863	0.891	0.896	0.9071
	8	0.692	0.721	0.756	0.785	0.788	0.816	0.835	0.865	0.871	0.884
	9	0.661	0.690	0.726	0.756	0.759	0.788	0.808	0.840	0.847	0.860
	100	0.633	0.662	0.698	0.728	0.732	0.761	0.782	0.816	0.823	0.837
100.....	1	0.9 <sup>2</sup> 83	0.9 <sup>2</sup> 90	0.9 <sup>3</sup> 49	0.9 <sup>3</sup> 75	0.9 <sup>3</sup> 77	0.9 <sup>4</sup> 00	0.9 <sup>4</sup> 50	0.9 <sup>4</sup> 87	0.9 <sup>5</sup> 01	0.9 <sup>5</sup> 50
	2	0.9 <sup>2</sup> 30	0.9 <sup>2</sup> 48	0.9 <sup>2</sup> 65	0.9 <sup>2</sup> 76	0.9 <sup>2</sup> 77	0.9 <sup>2</sup> 85	0.9 <sup>2</sup> 90	0.9 <sup>3</sup> 48	0.9 <sup>3</sup> 55	0.9 <sup>3</sup> 69
	3	0.9867	0.9893	0.9 <sup>2</sup> 20	0.9 <sup>2</sup> 40	0.9 <sup>2</sup> 42	0.9 <sup>2</sup> 57	0.9 <sup>2</sup> 67	0.9 <sup>2</sup> 79	0.9 <sup>2</sup> 81	0.9 <sup>2</sup> 85
	4	0.9799	0.9831	0.9868	0.9894	0.9897	0.9 <sup>2</sup> 20	0.9 <sup>2</sup> 35	0.9 <sup>2</sup> 55	0.9 <sup>2</sup> 58	0.9 <sup>2</sup> 65
	5	0.9728	0.9767	0.9811	0.9844	0.9847	0.9877	0.9896	0.9 <sup>2</sup> 23	0.9 <sup>2</sup> 28	0.9 <sup>2</sup> 39
	6	0.9656	0.9700	0.9751	0.9789	0.9794	0.9829	0.9853	0.9887	0.9894	0.9 <sup>2</sup> 07
	7	0.9584	0.9632	0.9689	0.9733	0.9738	0.9778	0.9806	0.9847	0.9855	0.9871
	8	0.9512	0.9564	0.9626	0.9675	0.9681	0.9726	0.9757	0.9804	0.9813	0.9832
	9	0.9439	0.9496	0.9562	0.9615	0.9622	0.9671	0.9706	0.9759	0.9769	0.9791
	100	0.9367	0.9427	0.9498	0.9555	0.9562	0.9616	0.9654	0.9712	0.9723	0.9747
	100	0.538	0.548	0.560	0.571	0.573	0.584	0.593	0.607	0.610	0.617

<sup>a</sup> Upper limit for ratio type 1 event to type 1 + type 2. Calculate lower limit from this table with lower limit  $(n_1, n_2) = 1 -$  upper limit  $(n_2, n_1)$ .

<sup>b</sup> 0.9<sup>2</sup>00 means 0.9900; and in general for 9<sup>n</sup>.

any specific values of  $n_1$ ,  $n_2$ , and CL, exact numerical solutions can be computed to arbitrary accuracy.

Table 6 lists exact numerically determined values for  $p_{1u}$  at the previously chosen confidence levels for a number of values of  $n_1$  and  $n_2$ . A separate table is not required for the lower limits because they can be simply obtained from the upper limits. The relationship between the two, derived from equations (16) and (17), is

$$p_{1l} = 1 - p_{2u}, \quad (18)$$

where  $p_{2u}$  is the upper limit for the ratio of type 2 events to total events and can be obtained from Table 6 by switching  $n_1$  and  $n_2$ . For example, if the observed number of events are  $n_1 = 6$  and  $n_2 = 4$ , the best guess for the ratio of type 1 to total is 0.6, and the 99% confidence lower limit is 1 minus the upper limit from the table for  $n_1 = 4$ ,  $n_2 = 6$ , or  $1 - 0.782 = 0.218$ . The 99% confidence upper limit is 0.907 directly from Table 6, and the 99% confidence double-sided interval (CL = 0.995 column) is 0.191–0.923.

In many cases, the quantity of interest is the ratio of type 1 to



type 2 event,  $r_{12}$ , instead of type 1 to total. The upper and lower limits for  $r_{12}$  are related to  $p_{1u}$  and  $p_{1l}$  by

$$r_{12u} = \frac{p_{1u}}{1 - p_{1u}}, \quad (19)$$

$$r_{12l} = \frac{p_{1l}}{1 - p_{1l}} = \frac{1 - p_{2u}}{p_{2u}}. \quad (20)$$

Using the above example of  $n_1 = 6$  and  $n_2 = 4$ , the best guess for  $r_{12}$  is 1.5, and the 99% confidence upper and lower limits (single-sided) are 9.73 and 0.279.

### c) Algebraic Expressions

There are two special cases where exact expressions for  $p_{1u}$  can be derived from equation (16). These are

$$p_{1u} = \begin{cases} \text{CL}^{1/(n_1+n_2)} & (n_2 = 1) \\ 1 - (1 - \text{CL})^{1/n_2} & (n_1 = 0) \end{cases}. \quad (21)$$

Lower limits for  $n_1 = 1$  and  $n_2 = 0$  can be obtained from these expressions using equation (18). For instance,  $p_{1l}(n_2 = 0) = (1 - \text{CL})^{1/n_1}$ .

For other values of  $n_1$  and  $n_2$ , we make use of the relation between the binomial sum and the incomplete beta function,  $I(p|\alpha, \beta) \equiv I_p(\alpha, \beta)$ ,

$$\sum_{x=\alpha}^n \binom{n}{x} p^x (1-p)^{n-x} = I(p|\alpha, n-\alpha+1) \quad (22)$$

(see, e.g., Abramowitz and Stegun 1965). Combining this relation with equations (16) and (17) gives

$$I(p_{1u}|n_1+1, n_2) = \text{CL}, \quad (23)$$

$$I(p_{1l}|n_1, n_2+1) = 1 - \text{CL}. \quad (24)$$

Thus,  $p_{1u}$  and  $p_{1l}$  can be obtained using tables of percentage points of the incomplete  $\beta$ -function (see, e.g., Beyer 1966) in the same manner that  $\chi^2$  tables are used to give Poisson limits. As with the Poisson limits, the technique is straightforward but cumbersome since several algebraic steps are involved and, in many cases, interpolations in the tables are required.

Equation (23) is useful for developing an approximate expression for  $p_{1u}$ . An approximate inverse function for the incomplete  $\beta$ -function is given by Abramowitz and Stegun (1965) as follows: if  $I(x_a|\alpha, \beta) = 1 - a$  and  $y_a$  is defined such that  $G(y_a) = a$  [see § IIb(i)], then

$$x_a \approx \frac{\alpha}{\alpha + \beta e^{2y_a}}, \quad (25)$$

where

$$w = \frac{y_a(h+\lambda)^{1/2}}{h} - \left( \frac{1}{2\beta-1} - \frac{1}{2\alpha-1} \right) \left( \lambda + \frac{5}{6} - \frac{2}{3h} \right),$$

$$h = 2 \left( \frac{1}{2\beta-1} + \frac{1}{2\alpha-1} \right)^{-1},$$

and

$$\lambda = \frac{y_a^2 - 3}{6}.$$

Combining equations (23) and (25), adding a correction term, and using the variable  $S$  defined in § IIb(i) (see Table 3), we derive an approximate expression for  $p_{1u}$ ,

$$p_{1u} \approx \frac{(n_1+1)e^{2w} + \epsilon n_2}{(n_1+1)e^{2w} + n_2}, \quad (26)$$

where

$$w = \frac{S(h+\lambda)^{1/2}}{h} + \left( \frac{1}{2n_2-1} - \frac{1}{2n_1+1} \right) \left( \lambda + \frac{5}{6} - \frac{2}{3h} \right),$$

$$h = 2 \left( \frac{1}{2n_2-1} + \frac{1}{2n_1+1} \right)^{-1},$$

$$\lambda = \frac{S^2 - 3}{6},$$

and

$$\epsilon = 0.64(1-S)e^{-n_2}.$$

I was not able to find a simpler expression than equation (26) that is generally useful for all confidence levels and all values of  $n_1$  and  $n_2$ . For cases where equations (21) do not apply (i.e., for  $n_2 \geq 2$  and  $n_1 \geq 1$ ), equation (26) gives upper limits accurate to better than 4% for all confidence levels. If  $n_1 \geq 4$  and  $n_2 \geq 4$ , equation (26) is accurate to better than 1%. These two percentage accuracies also hold for  $1 - p_{1u}$ , which is the quantity used to calculate lower limits. In the Gaussian limit of large  $n_1$  and  $n_2$ , the upper limit is given by

$$p_{1u} = \frac{n_1}{n_1 + n_2} + \frac{S\sqrt{n_1 n_2}}{(n_1 + n_2)^{3/2}}. \quad (27)$$

This expression becomes accurate at the 1% level only for  $n_1$  and  $n_2$  both greater than  $\sim 50$ .

*Recommended expressions:* For  $n_1 = 0$  or  $n_2 = 1$ , use the exact equations (21) for the binomial upper limits. For other values of  $n_1$  and  $n_2$  use equation (26), which gives 4% accuracy. Calculate the lower limits from these upper limit expressions using  $p_{1l} = 1 - p_{2u}$ , where  $p_{2u}$  is the upper limit with  $n_1$  and  $n_2$  switched. This relationship combined with equations (21) gives exact lower limits for  $n_1 = 1$  or  $n_2 = 0$  and combined with equation (26) gives lower limits accurate to within 4% for all other values.

## IV. SUMMARY

Poisson statistics apply when event rates are calculated from small numbers of observed events. For  $n$  observed events, the single-sided upper limit,  $\lambda_u$ , and single-sided lower limit,  $\lambda_l$ , of confidence level CL are given by equations (1) and (2). Previous techniques for obtaining actual limit values from these equations were cumbersome. This paper has presented tables of upper and lower limits (Tables 1 and 2) for  $n = 0-50$  for all common confidence levels used in astrophysics. Also, convenient approximate expressions for  $\lambda_u$  and  $\lambda_l$  were developed, the most useful of which are  $\lambda_u \approx n + S(n+1)^{1/2} + (S^2+2)/3$  (good to better than 10% for all  $n$  and CL) and better than to 1% for  $n \geq 4$  and  $\lambda_l \approx n[1 - 1/9n - S/3(n)^{1/2} + \beta n^2]^3$  (good to better than 2% for all  $n$  and CL), where  $S$ ,  $\beta$ , and  $\gamma$  are given in Table 3.

Binomial statistics apply when ratios of rates of two different event types are calculated from small numbers of observed events. For  $n_1$  observed events of type 1 and  $n_2$  of type 2, the single-sided upper limit,  $p_{1u}$ , and single-sided lower limit,  $p_{1l}$ , of confidence level CL to the ratio of type 1 event rate to type 1 plus type 2 are given by equations (16) and (17). Again, in this case, previous techniques for obtaining values from these equations were cumbersome. Considering upper limits first, Table 6 lists values for  $p_{1u}$  for a number of small  $n_1$  and  $n_2$  for all common confidence levels used in astrophysics. There are two special cases with exact expressions for the upper limits:  $p_{1u} =$

$CL^{1/n_1+n_2}$  ( $n_2 = 1$ ) and  $p_{1u} = 1 - (1 - CL)^{1/n_2}$  ( $n_1 = 0$ ). The approximate expression in equation (26) gives values for  $p_{1u}$  accurate to better than 4% for all other  $n_1$  and  $n_2$ , and to better than 1% for  $n_1 \geq 4$  and  $n_2 \geq 4$ . Lower limits can be obtained directly from the upper limits in the table or in the above equations by taking 1 minus the upper limit value, for  $n_1$  and  $n_2$  exchanged.

I greatly appreciate the comments and advice of Martin H. Israel. This paper is a development from an unpublished report by Dr. Israel, in which equations (7) and (11) were derived. Thanks are also due to William A. Wheaton for useful discussions concerning Poisson statistics. I am grateful for E. Schronce for typing the manuscript of this paper.

## APPENDIX

### NUMERICAL VERIFICATION OF POISSON LIMIT EQUATIONS

The purpose of this appendix is to demonstrate that the Poisson upper and lower limits defined by equations (1) and (2) in § II satisfy the definition of confidence limits and do so in an optimum manner. To simplify the discussion we will first consider only upper limits and then generalize the results to include lower limits.

For a quantity with a continuous distribution function such as a Gaussian distribution, upper limits are defined to be confidence limits of confidence level CL if they satisfy the following condition; for a large number of observers measuring a given physical quantity (with measurements distributed about the true value according to the distribution function) and each observer assigning an upper limit based on his or her measured value,  $100 \times CL\%$  of them will have upper limits greater than the true value (see, e.g., Cramér 1945). For a quantity with a discrete distribution function, such as a Poisson distribution, it is not possible to generate upper limits that satisfy this condition for all parameter values, so the definition becomes: *at least*  $100 \times CL\%$  of the observers have upper limits greater than the true value. The lower limit condition is that at least  $100 \times CL\%$  of the observers have lower limits less than the true value.

Calculations have been performed to verify that the Poisson upper and lower limits in equations (1) and (2) satisfy this definition of confidence limits. A finely spaced sample of assumed true event rates,  $\lambda_t$ , was chosen. For each  $\lambda_t$ , a distribution of observed values was then determined from the Poisson distribution function

$$P(n) = \frac{\lambda_t^n e^{-\lambda_t}}{n!}. \quad (\text{A1})$$

For each observed value, upper and lower limits of confidence level 0.8413 (corresponding to  $1 \sigma$  Gaussian limits) were obtained from Tables 1 and 2. A tally was then taken of all observed values to determine the fraction of observers with upper limits greater than the true rate and the fraction with lower limits less than the true rate. The results are plotted as functions of true rate in Figure 2. As expected for a discrete distribution, the fractions vary as functions of true rate, dipping to exactly 0.8413 each time the true rate




FIG. 2.—Fraction of observers with (a) upper limit greater than the true rate  $\lambda_t$ , and (b) lower limit less than  $\lambda_t$ , as functions of  $\lambda_t$ . For all values of  $\lambda_t$ , the fraction is greater than or equal to the confidence level of 0.8413.

passes one of the limit values. The size of the variation decreases with increasing  $\lambda_i$ , approaching a constant value of 0.8413 in the limit. This is as expected since the discrete Poisson distribution approaches the continuous Gaussian distribution in the limit of large  $\lambda_i$ . The confidence limit definitions are satisfied for all rates; at least 84.13% of the observers have upper limits greater than the true rate and lower limits less than the true rate, for all  $\lambda_i$ . The limits are optimum in the sense that the fractions are exactly 0.8413 at some discrete values of the true rate. If any one of the upper limits in Table 1 were decreased or lower limits in Table 2 increased, the fraction would dip below 0.8413 for some true rate.

## REFERENCES

- Abramowitz, M., and Stegun, I. A. eds. 1965, *Handbook of Mathematical Functions* (New York: Dover).
- Beyer, W. H., ed. 1966, *CRC Handbook of Tables for Probability and Statistics* (2d ed.; Boca Raton: CRC Press).
- Cramér, H. 1945, *Mathematical Methods of Statistics* (Princeton: Princeton University Press).
- Israel, M. H. 1968, Caltech Space Radiation Lab. Internal Rept., No. 4.
- Pearson, E. S., and Hartley, H. O., eds. 1966, *Biometrika Tables for Statisticians*, Vol. 1 (3d ed.; Cambridge: Cambridge University Press).

NEIL GEHRELS: Code 661, Laboratory for High Energy Astrophysics, NASA/Goddard Space Flight Center, Greenbelt, MD 20771